

NOTES ON *PARABLASTOTHRIX NEARCTICA*
(HYMENOPTERA: ENCYRTIDAE)

ROBERT L. ZUPARKO

Laboratory of Biological Control, University of California, Berkeley, 1050 San Pablo Avenue, Albany, California 94706.

Abstract.—New host records and a range extension are reported for *Parablastothrix nearctica*, as well as inaccuracies in the original description. Only one adult emerges from each host, suggesting the species is monoembryonic. *Parablastothrix* has been placed in the Copidosomatini, but its mode of reproduction contrasts with the other genera of the tribe, all of which appear to consist of polyembryonic species.

Key Words: *Parablastothrix nearctica*, *Paradelius rubra*, *Stigmella variella*, Copidosomatini, polyembryony

Parablastothrix nearctica Miller (Hymenoptera: Encyrtidae) was described from specimens collected in the eastern United States and reared from two leafminers, a *Coptodisca* sp. (Lepidoptera: Heliozelidae) and an *Obrussa* sp. (Lepidoptera: Nepticulidae) (Miller 1965). In parts of this and some subsequent papers, this species was also referred to as *P. nearcticus*: this suffix is incorrect because the generic name is based on the Greek feminine *thrix*.

In March 1994, I collected five cocoons from the underside of leaves of a coast live oak (*Quercus agrifolia* Nee) at our experiment station in Albany, California. Three of these were formed by *Stigmella variella* (Braun) (Lepidoptera: Nepticulidae), from each of which emerged a single *Parablastothrix* adult (two females and one male). A male *Paradelius rubra* Whitfield (Hymenoptera: Braconidae) emerged from the fourth cocoon.

The *Parablastothrix* adults exhibited a frenetic activity, typical of many hyperparasitoids. But when I dissected the four empty cocoons, I found no more than one meconial pellet in each cocoon, indicating both

species were solitary, primary parasitoids. The fifth cocoon held a dead, well-formed *P. rubra* male pupa. In October 1994, I collected 10 more *Parablastothrix* adults (nine females and one male) from the foliage of the same trees.

The *Parablastothrix* specimens were identical to those reported in a study on parasitoids of leafmining lepidopterans on oaks in northern California, dating back to 1961 (Green 1979). All specimens had yellow legs with darkened metatibiae, and a green body with a bronze hue, but which otherwise matched the description of *P. nearctica*. Green (1979) reported this encyrtid emerged individually from cocoons of *S. variella*, and from undetermined stages of *Coptodisca powellella* Opler, *Bucculatrix albertiella* Busck (Lepidoptera: Lyonetiidae) and *Phyllonorycter sandraella* (Opler) (Lepidoptera: Gracillariidae), while still in their leaf mines.

Miller (1965) reported that *P. nearctica* had white legs, and the overall body color was black with bluish and greenish reflections. Reexamination of the holotype and two paratypes of *P. nearctica* revealed that

the original description was in error. All three type specimens have dark metatibiae and a green body, identical to the California specimens. Thus the latter represent a range extension of *P. nearctica*. I have seen additional specimens of *P. nearctica* collected from Arizona and central and southern California.

Hosts have been recorded for only 8 of the 17 described *Parablastothrix* species—all are leaf-mining Lepidoptera from the families Gracillariidae, Lyonetiidae, and Nepticulidae (Hedqvist 1976, Logvinovskaya 1981, Khan 1983, Trjapitzin 1989). An undescribed *Parablastothrix* sp. was reported as a solitary, primary parasitoid on a nepticulid, a heliozelid, and a gracillariid on pecan in Georgia (Dutcher and Heyerdahl 1988).

Hoffer (1955) placed *Parablastothrix* in the tribe Microteriyini (subtribe Pentacnemii), with *Calometopia* Mercet and *Pentacnemus* Howard. Noyes and Hayat (1984) synonymized *Pentacnemus* with *Copidosoma* Ratzeburg, and Trjapitzin (1989) synonymized *Calometopia* with *Parablastothrix*. Trjapitzin and Gordh (1978) placed *Parablastothrix* in the subtribe Parablastothrichina within the Copidosomatini, but remarked on the absence of proven polyembryony. Noyes and Hayat (1984) considered the genera in this subtribe to be morphologically closer to the Aphycina (tribe Aphycini); however they did not formally place these genera in their scheme of systematic relationships. Recent work by G. Zolnerowich (pers. comm.) retains *Parablastothrix* as a more basal lineage within the Copidosomatini.

Within the Copidosomatini, polyembryony appears to be the rule, having been reported in all species whose biologies have been investigated, including species of *Ap-silophrys* De Santis, *Copidosoma* (= *Litomastix* Thomson), *Copidosomopsis* Girault, *Paralitomastix* Mercet, *Ageniaspis* Dahlbom, *Holcothorax* Mayr and *Paraleurocerus* Girault (Noyes 1980, Noyes and Hayat 1984). Polyembryony may also occur in

Coelopencyrtus Timberlake (Timberlake 1919, Trjapitzin 1960, Taylor 1961, Annecke 1968), but this mode of reproduction has never been conclusively demonstrated.

Solitary parasitism presupposes monoembryony. However, such evidence is not conclusive, because the polyembryonic species *Macrocentrus ancyliivorus* Rohwer (Hymenoptera: Braconidae) is known to develop as a solitary parasitoid (Daniel 1932). Embryological studies have not been conducted on any *Parablastothrix* species. Nevertheless, the solitary habit of *P. nearctica* and the absence of any mention of multiple emergences from all other *Parablastothrix* rearing records, suggests that polyembryony is absent in the entire genus. Such a lifestyle would then biologically separate this genus from the remainder of the Copidosomatini.

Material examined.—ARIZONA. COCHISE Co.: 10 mi. W. Portal, on *Juniperus*. CALIFORNIA. ALAMEDA Co.: Albany, *Stigmella variella* on *Quercus agrifolia*; Berkeley Hills, *S. variella* on *Q. agrifolia* and *Phyllonorycter sandraella*; Patterson Reserve, Del Valle Lake, *Stigmella* sp.; CONTRA COSTA Co.: Antioch, *S. variella* on *Q. agrifolia*; 3 km E of Antioch, *S. variella* on *Q. agrifolia*; Cowell, *S. variella* on *Q. agrifolia*; Oakley, *S. variella* on *Q. agrifolia*; Russellmann Park; Russell Tree Farm, 6 km NE of Orinda, *S. variella*; MARIN Co.: Novato, *Coptodisca powellella* on *Q. agrifolia*; Sausalito, on *Q. agrifolia*; MONTEREY Co.: 10 km SE of Big Sur; SAN MATEO Co.: Woodside, *Bucculatrix albertiella* on *Q. agrifolia*; SANTA BARBARA Co.: Santa Cruz Island, Canada de la Cuesta, *Stigmella* sp. on *Q. dumosa*; SONOMA Co.: Sonoma, *Stigmella* sp. on *Q. agrifolia*. MICHIGAN. MIDLANDS Co.. VIRGINIA. Falls Church. MISSISSIPPI. HINDS Co.: Jackson, *Coptodisca* sp. on *Vaccinium arboreum* Marsh, holotype (no. 8800, deposited in Canadian National Collection, Ottawa, Ontario) and paratype (deposited in United States National Museum, Washington, D.C.). WEST VIRGINIA, MONON-

GALIA Co.: Morgantown, *Obrussa* sp., paratype (deposited in U.S.N.M.).

ACKNOWLEDGMENTS

I thank Jerry Powell, University of California, Berkeley, for identifying the *S. variegata* cocoons, and Michael Schauff (U.S.N.M.), Vince Lee (California Academy of Sciences) and Steven Heydon (U.C., Davis) for providing access to material. Gary Gibson of the Biosystematics Research Centre (Agriculture Canada) kindly reexamined the holotype of *P. nearctica*. Ken Hagen and Leo Caltagirone (U.C., Berkeley) and Gregory Zolnerowich (Texas A&M University) provided valuable discussions and advice.

LITERATURE CITED

- Annecke, D. P. 1968. Records and descriptions of African Encyrtidae 4 (Hymenoptera: Chalcidoidea). *Journal of the Entomological Society of South Africa* 31: 249-264.
- Daniel, D. M. 1932. *Macrocentrus ancylivorus* Rohwer, a polyembryonic braconid parasite of the Oriental fruit moth. New York State Agricultural Experiment Station, Technical Bulletin #187. 101 pp.
- Dutcher, J. D. and R. Heyerdahl. 1988. Parasitic Hymenoptera of the leafmining guild of pecan, pp. 445-458. In Gupta, V.K., ed., *Advances in Parasitic Hymenoptera Research*, E. J. Brill, Leiden.
- Green, D. S. 1979. Ecology and host-specificity of parasitoids of leaf-mining Lepidoptera on *Quercus agrifolia* (Fagaceae) in California. Ph.D. Thesis, University of California, Berkeley. 204 pp.
- Hedqvist, K.-J. 1976. Descriptions of new chalcid flies (Hym., Chalcidoidea, Encyrtidae and Eulophidae) reared from *Nepticula* species collected in Sri Lanka. *Entomologisk Tidskrift* 97: 50-54.
- Hoffer, A. 1955. The phylogeny and taxonomy of the family Encyrtidae (Hym., Chalcidoidea). *Sbornik Narodniho Museu v Praze* 11: 1-22.
- Khan, M. A. 1983. A new species of *Parablastothrix* (Hymenoptera: Encyrtidae) from India. *Journal, Bombay Natural History Society* 80: 180-182.
- Logvinovskaya, T. V. 1981. A review of Holarctic species of the genus *Parablastothrix* Mercet (Hymenoptera, Encyrtidae). *Entomological Review* 60: 147-153.
- Miller, C. D. F. 1965. A Nearctic species of *Parablastothrix* Mercet (Hymenoptera: Encyrtidae). *Canadian Entomologist* 97: 750-753.
- Noyes, J. S. 1980. A review of the genera of Neotropical Encyrtidae (Hymenoptera: Chalcidoidea). *Bulletin of the British Museum (Natural History)* 41: 107-253.
- Noyes, J. S. and M. Hayat. 1984. A review of the genera of Indo-Pacific Encyrtidae (Hymenoptera: Chalcidoidea). *Bulletin of the British Museum (Natural History)* 48: 131-395.
- Taylor, J. S. 1961. A note on some insects associated with Xylocopidae in the eastern Cape Province, South Africa. *Pan-Pacific Entomologist* 37: 220-222.
- Timberlake, P. H. 1919. Descriptions of new genera and species of Hawaiian Encyrtidae (Hymenoptera). *Proceedings of the Hawaiian Entomological Society* 4: 197-231.
- Trjapitzin, V. A. 1960. A Palaearctic species of the genus *Coelopencyrtus* Timb. (Hymenoptera Encyrtidae). *Entomological Review* 39: 500-502.
- . 1989. Parasitic Hymenoptera of the Family Encyrtidae of Palaearctics. Leningrad, "Nauka," Leningrad Division. 488 pp.
- Trjapitzin, V. A. and G. Gordh. 1978. A review of the Nearctic Encyrtidae (Hymenoptera, Chalcidoidea). *Communication II. Entomological Review* 57: 437-448.